arXiv:1508.06626v1 [math.CA] 26 Aug 2015

ON THE MAIN EQUATION OF INVERSE STURM-LIOUVILLE OPERATOR WITH DISCONTINUOUS COEFFICIENT

KH. R. MAMEDOV AND D. KARAHAN

Abstract. In this work, a boundary value problem for Sturm-Liouville operator with discontinuous coefficient is examined. The main equation is obtained which has an important role in solution of inverse problem for boundary value problem and uniqueness of its solution is proved. Uniqueness theorem for the solution of the inverse problem is given.

1. Introduction

In many practices, spectral problems are faced for differential equations which have discontinuous coefficient and discontinuity conditions in interval ([1]-[8]). These problems generally emerge in physics, mechanics and geophysics in non-homogeneous and discontinuous environments.

We consider a heat problem in a rod which is composed of materials having different densities. In the initial time, let the temperature is given arbitrary. Let be the temperature is zero in one end of the rod and the heat is isolated at the other end of the rod. In this case the heat flow in non-homogeneous rod is expressed with the following boundary problem:

$$\rho\left(x\right)\frac{\partial u}{\partial t}=\frac{\partial^{2}u}{\partial x^{2}}+q\left(x\right)u,\qquad 0<x<\pi,\ t>0,$$
$$\left.\frac{\partial u}{\partial x}\right|_{x=0}=0,\qquad u|_{x=\pi}=0,\quad t>0,$$

where $\rho\left(x\right)$, $q\left(x\right)$ are physical parameters and have specific properties. For instance, $\rho\left(x\right)$ defines the density of the material and piecewise-continuous function. Applying the method of separation of variables to this problem, we get the spectral problem below:

$$-y''+q(x)y=\lambda^{2}\rho(x)y,\ 0\leq x\leq\pi \tag{1.1}$$

$$y'(0)=y(\pi)=0, \tag{1.2}$$

here $q\left(x\right)\in L_{2}\left(0,\pi\right)$ is a real-valued function, $\rho\left(x\right)$ piecewise-continuous function the following:

$$\rho(x)=\begin{cases}1, & 0\leq x\leq a,\\ \alpha^{2}, & a<x\leq\pi\end{cases} \tag{1.3}$$

2010 *Mathematics Subject Classification.* 34A55; 34K10.

Key words and phrases. main equation; discontinuous Sturm-Liouville operator; inverse problem; boundary value problem; spectral analysis.

λ is spectral parameter and $a(1+\alpha) > \pi\alpha$.

When $\rho(x) \equiv 1$ or $\alpha = 1$, that is, in continuous case, the solution of inverse problem is given in [9]-[19]. The spectral properties of Sturm-Liouville opeator with discontinuos coefficient in different boundary conditions are examined in [20]-[23].

In this study, the main equation is obtained which has an important role in solution of inverse problem for boundary value problem and according to spectral data, the uniqueness of solution of inverse problem is proved. Similar problems are examined for the equation (1.1) with different boundary conditions in [24].

It was proved (see [25]), that the solution $\varphi(x,\lambda)$ of the equation (1.1) with initial conditions $\varphi(0,\lambda) = 1$, $\varphi'(0,\lambda) = 0$ can be represented as

$$\varphi(x,\lambda) = \varphi_0(x,\lambda) + \int_0^{\mu^+(x)} A(x,t)\cos\lambda t dt, \tag{1.4}$$

where $A(x,t)$ belongs to the space $L_{2,\rho}(0,\pi)$ for each fixed $x \in [0,\pi]$ and is related with the coefficient $q(x)$ of the equation (1.1) by the formula:

$$\frac{d}{dx}A(x,\mu^+(x)) = \frac{1}{4\sqrt{\rho(x)}}\left(1+\frac{1}{\sqrt{\rho(x)}}\right)q(x), \tag{1.5}$$

$$\varphi_0(x,\lambda) = \frac{1}{2}\left(1+\frac{1}{\sqrt{\rho(x)}}\right)\cos\lambda\mu^+(x) + \frac{1}{2}\left(1-\frac{1}{\sqrt{\rho(x)}}\right)\cos\lambda\mu^-(x) \tag{1.6}$$

is the solution of (1.1) when $q(x) \equiv 0$,

$$\mu^+(x) = \pm x\sqrt{\rho(x)} + a\left(1 \mp \sqrt{\rho(x)}\right) \tag{1.7}$$

It is similarly shown in [24], [21] that the roots of the equation $\varphi(\pi,\lambda) = 0$ have the following form

$$\lambda_n = \lambda_n^0 + \frac{d_n}{\lambda_n^0} + \frac{k_n}{n},\ \lambda_n \geq 0,$$

where $\left\{\lambda_n^0\right\}^2$ are the eigenvalues of problem (1.1), (1.2) when $q(x) \equiv 0$, d_n is a bounded sequence, $k_n \in l_2$ and norming constants:

$$\alpha_n = \int_0^\pi \rho(x)\varphi^2(x,\lambda_n)dx.$$

2. Main Equation

Theorem 2.1. *For each fixed $x \in [0,\pi]$ the kernel $A(x,t)$ from the representation (1.4) satisfies the following linear functional integral equation*

$$\frac{2}{1+\sqrt{\rho(t)}}A\left(x,\mu^+(t)\right) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}}A\left(x,2a-t\right) +$$

$$+ F(x,t) + \int_0^{\mu^+(x)} A(x,\xi)F_0(\xi,t)d\xi = 0, \quad 0 < t < x \tag{2.1}$$

where

$$F_0(x,t) = \sum_{n=1}^{\infty}\left(\frac{\varphi_0(t,\lambda_n)\cos\lambda_n x}{\alpha_n} - \frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0 x}{\alpha_n^0}\right) \tag{2.2}$$

$$F(x,t)=\frac{1}{2}\left(1+\frac{1}{\sqrt{\rho(x)}}\right)F_0(\mu^+(x),t)+\frac{1}{2}\left(1-\frac{1}{\sqrt{\rho(x)}}\right)F_0(\mu^-(x),t) \tag{2.3}$$

$\left\{\lambda_n^0\right\}^2$ are eigenvalues and α_n^0 are norming constants of the boundary value problem (1.1), (1.2) when $q(x)\equiv 0$.

Proof. From (1.4) we have

$$\varphi_0(x,\lambda)=\varphi(x,\lambda)-\int_0^{\mu^+(x)}A(x,t)\cos\lambda t dt. \tag{2.4}$$

It follows from (1.4) and (2.4) that

$$\sum_{n=1}^N\frac{\varphi(x,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}=\sum_{n=1}^N\left(\frac{\varphi_0(x,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}+\right.$$
$$\left.+\frac{\varphi_0(t,\lambda_n)}{\alpha_n}\int_0^{\mu^+(x)}A(x,\xi)\cos\lambda_n\xi d\xi\right)=$$
$$=\sum_{n=1}^N\left(\frac{\varphi_0(x,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}-\frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}\right)+$$
$$+\sum_{n=1}^N\frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}+$$
$$+\int_0^{\mu^+(x)}A(x,\xi)\sum_{n=1}^N\left(\frac{\varphi_0(t,\lambda_n)\cos\lambda_n\xi}{\alpha_n}-\frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0}\right)d\xi+$$
$$+\int_0^{\mu^+(x)}A(x,\xi)\sum_{n=1}^N\frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0}d\xi$$
$$\sum_{n=1}^N\frac{\varphi(x,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}=\sum_{n=1}^N\frac{\varphi(x,\lambda_n)\varphi(t,\lambda_n)}{\alpha_n}-$$
$$-\int_0^{\mu^+(t)}A(t,\xi)\sum_{n=1}^N\frac{\varphi(x,\lambda_n)\cos\lambda_n\xi}{\alpha_n}d\xi.$$

Using the last two equalities, we obtain

$$\sum_{n=1}^N\left(\frac{\varphi(x,\lambda_n)\varphi(t,\lambda_n)}{\alpha_n}-\frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}\right)=$$
$$=\sum_{n=1}^N\left(\frac{\varphi_0(x,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}-\frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}\right)+$$
$$+\int_0^{\mu^+(x)}A(x,\xi)\sum_{n=1}^N\frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0}d\xi+$$
$$+\int_0^{\mu^+(x)}A(x,\xi)\sum_{n=1}^N\left(\frac{\varphi_0(t,\lambda_n)\cos\lambda_n\xi}{\alpha_n}-\frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0}\right)d\xi+$$
$$+\int_0^{\mu^+(t)}A(t,\xi)\sum_{n=1}^N\frac{\varphi(x,\lambda_n)\cos\lambda_n\xi}{\alpha_n}d\xi,$$

or

$$\Phi_N(x,t) = I_{N1}(x,t) + I_{N2}(x,t) + I_{N3}(x,t) + I_{N4}(x,t), \tag{2.5}$$

where

$$\Phi_N(x,t) := \sum_{n=1}^{N} \left(\frac{\varphi(x,\lambda_n)\varphi(t,\lambda_n)}{\alpha_n} - \frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0} \right),$$

$$I_{N1}(x,t) := \sum_{n=1}^{N} \left(\frac{\varphi_0(x,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n} - \frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0} \right),$$

$$I_{N2}(x,t) := \int_0^{\mu^+(x)} A(x,\xi) \sum_{n=1}^{N} \frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0} d\xi,$$

$$I_{N3}(x,t) := \int_0^{\mu^+(x)} A(x,\xi) \sum_{n=1}^{N} \left(\frac{\varphi_0(t,\lambda_n)\cos\lambda_n\xi}{\alpha_n} - \frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0} \right) d\xi,$$

$$I_{N4}(x,t) := \int_0^{\mu^+(t)} A(t,\xi) \sum_{n=1}^{N} \frac{\varphi(x,\lambda_n)\cos\lambda_n\xi}{\alpha_n} d\xi.$$

It is easily found by using (2.2) and (2.3)

$$F(x,t) = \sum_{n=1}^{\infty} \left(\frac{\varphi_0(x,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n} - \frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0} \right).$$

Let $f(x)$ be an absolutely continuous function, $f'(0) = f(\pi) = 0$. Then using expansion formula (see [21]),

$$\sum_{n=1}^{\infty} \int_0^{\pi} f(t)\rho(t) \frac{\varphi(x,\lambda_n)\varphi(t,\lambda_n)}{\alpha_n} dt = f(x),$$

$$\sum_{n=1}^{\infty} \int_0^{\pi} f(t)\rho(t) \frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0} dt = f(x). \tag{2.6}$$

Using (2.6) we have:

$$\lim_{N\to\infty} \max_{0\le x\le\pi} \left| \int_0^{\pi} f(t)\rho(t)\Phi_N(x,t)dt \right| =$$

$$= \lim_{N\to\infty} \max_{0\le x\le\pi} \left| \int_0^{\pi} f(t)\rho(t) \sum_{n=1}^{N} \left(\frac{\varphi(x,\lambda_n)\varphi(t,\lambda_n)}{\alpha_n} - \frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0} \right) dt \right| \le$$

$$\le \lim_{N\to\infty} \left\{ \max_{0\le x\le\pi} \left| \int_0^{\pi} f(t)\rho(t) \sum_{n=1}^{N} \frac{\varphi(x,\lambda_n)\varphi(t,\lambda_n)}{\alpha_n} dt - f(x) \right| + \right.$$

$$\left. + \max_{0\le x\le\pi} \left| \int_0^{\pi} f(t)\rho(t) \sum_{n=1}^{N} \frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0} dt - f(x) \right| \right\} = 0. \tag{2.7}$$

We obtain uniformly on $x \in [0,\pi]$

$$\lim_{N\to\infty} \int_0^{\pi} f(t)\rho(t) I_{N1}(x,t)dt =$$

$$= \lim_{N\to\infty}\int_0^\pi f(t)\rho(t)\sum_{n=1}^N\left(\frac{\varphi_0(x,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}-\frac{\varphi_0(x,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}\right)dt =$$

$$= \int_0^\pi f(t)\rho(t)F(x,t)dt. \tag{2.8}$$

It follows from (1.6) that

$$\cos\lambda\xi = \begin{cases} \varphi_0(\xi,\lambda) \quad , \quad \xi<a, \\ \frac{2\alpha}{1+\alpha}\varphi_0(\frac{\xi}{\alpha}+a-\frac{a}{\alpha},\lambda)+\frac{1-\alpha}{1+\alpha}\varphi_0(2a-\xi,\lambda) \quad , \quad \xi>a. \end{cases} \tag{2.9}$$

Taking into account (2.9) and (2.6), we get

$$\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)I_{N2}(x,t)dt =$$

$$= \lim_{N\to\infty}\int_0^\pi f(t)\rho(t)\int_0^{\mu^+(x)}A(x,\xi)\sum_{n=1}^N\frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0}d\xi dt =$$

$$= \lim_{N\to\infty}\int_0^\pi f(t)\rho(t)\int_0^a A(x,\xi)\sum_{n=1}^N\frac{\varphi_0(t,\lambda_n^0)\varphi_0(\xi,\lambda)}{\alpha_n^0}d\xi dt+$$

$$+\frac{2\alpha}{1+\alpha}\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)\int_a^{\alpha x-\alpha a+a}A(x,\xi)\times$$

$$\times\sum_{n=1}^N\frac{\varphi_0(t,\lambda_n^0)\varphi_0(\frac{\xi}{\alpha}+a-\frac{a}{\alpha},\lambda_n^0)}{\alpha_n^0}d\xi dt+$$

$$+\frac{1-\alpha}{1+\alpha}\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)\int_a^{\alpha x-\alpha a+a}A(x,\xi)\times$$

$$\times\sum_{n=1}^N\frac{\varphi_0(t,\lambda_n^0)\varphi_0(2a-\xi,\lambda_n^0)}{\alpha_n^0}d\xi dt =$$

$$= \int_0^a A(x,\xi)\int_0^\pi f(t)\rho(t)\sum_{n=1}^\infty\frac{\varphi_0(t,\lambda_n^0)\varphi_0(\xi,\lambda_n^0)}{\alpha_n^0}dtd\xi+$$

$$+\frac{2\alpha}{1+\alpha}\int_a^{\alpha x-\alpha a+a}A(x,\xi)\int_0^\pi f(t)\rho(t)\times$$

$$\times\sum_{n=1}^\infty\frac{\varphi_0(t,\lambda_n^0)\varphi_0(\frac{\xi}{\alpha}+a-\frac{a}{\alpha},\lambda_n^0)}{\alpha_n^0}dtd\xi+$$

$$+\frac{1-\alpha}{1+\alpha}\int_a^{\alpha x-\alpha a+a}A(x,\xi)\int_0^\pi f(t)\rho(t)\times$$

$$\times\sum_{n=1}^\infty\frac{\varphi_0(t,\lambda_n^0)\varphi_0(2a-\xi,\lambda_n^0)}{\alpha_n^0}dtd\xi =$$

$$= \int_0^a A(x,\xi)f(\xi)d\xi+\frac{2\alpha}{1+\alpha}\int_a^{\alpha x-\alpha a+a}A(x,\xi)f\left(\frac{\xi}{\alpha}+a-\frac{a}{\alpha}\right)d\xi+$$

$$+\frac{1-\alpha}{1+\alpha}\int_a^{\alpha x-\alpha a+a}A(x,\xi)f\left(2a-\xi\right)d\xi.$$

Substituting $\frac{\xi}{\alpha}+a-\frac{a}{\alpha}\to\xi'$ and $2a-\xi\to\xi''$ we obtain

$$\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)I_{N2}(x,t)dt=\int_0^a A(x,\xi)f\left(\xi\right)d\xi+$$

$$+\frac{2\alpha^2}{1+\alpha}\int_a^x A\left(x,\alpha\xi'-\alpha a+a\right)f\left(\xi'\right)d\xi'+$$

$$+\frac{1-\alpha}{1+\alpha}\int_{-\alpha x+\alpha a+a}^{a} A(x,2a-\xi'')f\left(\xi''\right)d\xi''.$$

Since $A(x,2a-\xi'')\equiv 0$ when $2a-\xi>\alpha x-\alpha a+a$, we have

$$\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)I_{N2}(x,t)dt=\int_0^a A(x,t)f\left(t\right)dt+$$

$$+\frac{2\alpha^2}{1+\alpha}\int_a^x A\left(x,\alpha t-\alpha a+a\right)f\left(t\right)dt+$$

$$+\frac{1-\alpha}{1+\alpha}\int_0^a A(x,2a-t)f\left(t\right)dt.$$

Thus, uniformly on $x\in[0,\pi]$:

$$\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)I_{N2}(x,t)dt=\int_0^x\frac{2\rho\left(t\right)}{1+\sqrt{\rho\left(t\right)}}A(x,\mu^+(t))f\left(t\right)dt+$$

$$+\int_0^x\frac{1-\sqrt{\rho\left(2a-t\right)}}{1+\sqrt{\rho\left(2a-t\right)}}A(x,2a-t)f\left(t\right)dt. \tag{2.10}$$

Using (2.2), uniformly on $x\in[0,\pi]$

$$\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)I_{N3}(x,t)dt=$$

$$=\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)\int_0^{\mu^+(x)}A(x,\xi)\sum_{n=1}^{N}\left(\frac{\varphi_0(t,\lambda_n)\cos\lambda_n\xi}{\alpha_n}-\right.$$

$$\left.-\frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0}\right)d\xi dt=$$

$$=\int_0^\pi f(t)\rho(t)\int_0^{\mu^+(x)}A(x,\xi)\sum_{n=1}^{\infty}\left(\frac{\varphi_0(t,\lambda_n)\cos\lambda_n\xi}{\alpha_n}-\right.$$

$$\left.-\frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0}\right)d\xi dt=$$

$$=\int_0^\pi f(t)\rho(t)\int_0^{\mu^+(x)}A(x,\xi)F_0(\xi,t)d\xi dt. \tag{2.11}$$

Using the residue theorem and the formula $\frac{\varphi(x,\lambda_n)}{2\lambda_n\alpha_n}=\frac{\psi(x,\lambda_n)}{\dot{\Delta}(\lambda_n)}$ (see [21]), where $\psi\left(x,\lambda\right)$ is the solution of (1.1) with initial condition $\psi\left(\pi,\lambda\right)=0$, $\psi'\left(\pi,\lambda\right)=1$ and $\Delta\left(\lambda\right)=\varphi\left(\pi,\lambda\right)$ is the characteristic function of (1.1)-(1.3), $\dot{\Delta}(\lambda)=\frac{d}{d\lambda}\Delta\left(\lambda\right)$, we calculate

$$\lim_{N\to\infty}\int_0^\pi f(t)\rho(t)I_{N4}(x,t)dt=$$

$$
\begin{aligned}
&= \lim_{N\to\infty}\int_0^{\pi} f(t)\rho(t)\int_0^{\mu^+(t)} A(t,\xi)\sum_{n=1}^{N}\frac{\varphi(x,\lambda_n)\cos\lambda_n\xi}{\alpha_n}d\xi dt =\\
&= 2\lim_{N\to\infty}\int_0^{\pi} f(t)\rho(t)\sum_{|\lambda_n|\le N}\lambda_n\frac{\psi\left(x,\lambda_n\right)}{\dot{\Delta}(\lambda_n)}\int_0^{\mu^+(t)} A(t,\xi)\cos\lambda_n\xi d\xi dt =\\
&= 2\lim_{N\to\infty}\int_0^{\pi} f(t)\rho(t)\sum_{|\lambda_n|\le N}\underset{\lambda=\lambda_n}{Res}\left[\lambda\frac{\psi\left(x,\lambda\right)}{\Delta(\lambda)}\int_0^{\mu^+(t)} A(t,\xi)\cos\lambda\xi d\xi\right]dt =\\
&= 2\lim_{N\to\infty}\int_0^{\pi} f(t)\rho(t)\frac{1}{2\pi i}\oint_{\Gamma_N}\lambda\frac{\psi\left(x,\lambda\right)}{\Delta(\lambda)}\int_0^{\mu^+(t)} A(t,\xi)\cos\lambda\xi d\xi d\lambda dt =\\
&= \lim_{N\to\infty}\int_0^{\pi} f(t)\rho(t)\frac{1}{\pi i}\oint_{\Gamma_N}\lambda\frac{\psi\left(x,\lambda\right)}{\Delta(\lambda)}e^{|Im\lambda|\mu^+(t)}e^{-|Im\lambda|\mu^+(t)}\times\\
&\qquad\times\int_0^{\mu^+(t)} A(t,\xi)\cos\lambda\xi d\xi d\lambda dt =\\
&= \int_0^{\pi} f(t)\rho(t)\lim_{N\to\infty}\left(\frac{1}{\pi i}\oint_{\Gamma_N}\lambda\frac{\psi\left(x,\lambda\right)}{\Delta(\lambda)}e^{|Im\lambda|\mu^+(t)}e^{-|Im\lambda|\mu^+(t)}\times\right.\\
&\qquad\left.\times\int_0^{\mu^+(t)} A(t,\xi)\cos\lambda\xi d\xi d\lambda\right)dt
\end{aligned}
\tag{2.12}
$$

where $\Gamma_N=\{\lambda:|\lambda|=N\}$. Since (see [21])

$$
\begin{aligned}
\psi(x,\lambda) &= O\left(\frac{e^{|Im\lambda|(\mu^+(\pi)-\mu^+(x))}}{|\lambda|}\right), \quad |\lambda|\to\infty,\\
|\Delta(\lambda)| &\ge C_\delta e^{|Im\lambda|\mu^+(\pi)}, \quad \lambda\in G_\delta,
\end{aligned}
$$

$(G_\delta=\{\lambda:|\lambda-\lambda_n|\ge\delta\}$, δ is a sufficiently small positive number) and according to Lemma 1.3.1 from [9]

$$
\lim_{|\lambda|\to\infty}\max_{0\le t\le\pi}e^{-|Im\lambda|\mu^+(t)}\left|\int_0^{\mu^+(t)}A(t,\xi)\cos\lambda\xi d\xi d\lambda\right|=0
$$

from the equality (2.12) we get

$$
\lim_{N\to\infty}\int_0^{\pi}f(t)\rho(t)I_{N4}(x,t)dt=0. \tag{2.13}
$$

Multiplying both sides of (2.5) by $\rho(x)f(x)$, integrating from 0 to π, tending to limit when $N\to\infty$ and using (2.7), (2.8), (2.10), (2.11) and (2.13) we have

$$
\int_0^x\frac{2\rho\left(t\right)}{1+\sqrt{\rho\left(t\right)}}A(x,\mu^+(t))f\left(t\right)dt+\int_0^x\frac{1-\sqrt{\rho\left(2a-t\right)}}{1+\sqrt{\rho\left(2a-t\right)}}A(x,2a-t)f\left(t\right)dt+
$$

$$
+\int_0^{\pi}f(t)\rho(t)F(x,t)dt+\int_0^{\pi}f(t)\rho(t)\int_0^{\mu^+(x)}A(x,\xi)F_0(\xi,t)d\xi dt=0.
$$

Since $f(x)$ can be chosen arbitrarily, we obtain

$$
\frac{2\rho\left(t\right)}{1+\sqrt{\rho\left(t\right)}}A(x,\mu^+(t))+\frac{1-\sqrt{\rho\left(2a-t\right)}}{1+\sqrt{\rho\left(2a-t\right)}}A(x,2a-t)+F(x,t)+
$$

$$+\int_0^{\mu^+(x)} A(x,\xi)F_0(\xi,t)d\xi = 0.$$

□

3. Theorem for the Solution of the Inverse Problem

Theorem 3.1. *For each fixed $x \in [0,\pi]$ main equation (2.1) has a unique solution $A(x,.) \in L_{2,\rho}\left(0,\mu^+(x)\right)$.*

Proof. We show that for each fixed $x > a$ the equation (2.1) is equivalent to the equation of the form $(I+B)\,f = g$ where B is a completely continuous operator, I is an identity operator in the space $L_{2,\rho}(0,\pi)$. (When $x \le a$ this fact is obvious.)

When $x > a$ rewrite (2.1) as

$$L_x A(x,.) + K_x A(x,.) = -F(x,.),$$

where

$$(L_x f)(t) = \begin{cases} f(t) + \frac{1-\alpha}{1+\alpha} f(2a-t) & , \quad t \le a < x, \\ \frac{2}{1+\alpha} f(\alpha t - \alpha a + a) & , \quad a < t < x. \end{cases} \tag{3.1}$$

$$(K_x f)(t) = \int_0^{\alpha x - \alpha a + a} f(\xi) F_0(\xi, t)\, d\xi, \quad 0 < t < x.$$

It is sufficient to prove that L_x is invertible, i.e. has a bounded inverse in $L_{2,\rho}\left(0,\pi\right)$.

Consider the equation $(L_x f)(t) = \phi(t)$, $\phi(t) \in L_{2,\rho}\left(0,\pi\right)$, i.e.

$$\begin{cases} f(t) + \frac{1-\alpha}{1+\alpha} f(2a-t) = \phi(t) & , \quad t \le a < x, \\ \frac{2}{1+\alpha} f(\alpha t - \alpha a + a) = \phi(t) & , \quad a < t < x. \end{cases}$$

From here it is easily to obtain

$$f(t) = \left(L_x^{-1}\phi\right)(t) = \begin{cases} \phi(t) - \frac{1-\alpha}{2}\phi\left(\frac{-t+\alpha a+a}{\alpha}\right) & , \quad t < a \\ \frac{1+\alpha}{2}\phi\left(\frac{t+\alpha a-a}{\alpha}\right) & , \quad t > a. \end{cases}$$

We show that

$$\|f\|_{L_2} = \left\|L_x^{-1}\phi\right\| \le C\,\|\phi\|_{L_2}.$$

In fact,

$$\int_0^\pi |f(t)|^2\,dt = \int_0^a \left|\phi(t) - \frac{1-\alpha}{2}\phi\left(\frac{-t+\alpha a+a}{\alpha}\right)\right|^2 dt +$$

$$+\int_a^\pi \left|\frac{1+\alpha}{2}\phi\left(\frac{t+\alpha a-a}{\alpha}\right)\right|^2 dt \le 2\int_0^a |\phi(t)|^2\,dt +$$

$$+2\left(\frac{1-\alpha}{2}\right)^2 \int_0^a \left|\phi\left(\frac{-t+\alpha a+a}{\alpha}\right)\right|^2 dt +$$

$$+\left(\frac{1+\alpha}{2}\right)^2 \int_a^\pi \left|\phi\left(\frac{t+\alpha a-a}{\alpha}\right)\right|^2 dt \le$$

$$\le 2\int_0^\pi |\phi(t)|^2\,dt + \frac{\alpha\,(1-\alpha)^2}{2}\int_a^{\frac{\alpha a+a}{\alpha}} |\phi(t)|^2\,dt +$$

$$+\alpha\left(\frac{1+\alpha}{2}\right)^2 \int_a^{\frac{\pi+\alpha a-a}{\alpha}} |\phi(t)|^2\,dt.$$

We put $\phi(t) = 0$, when $t > \pi$. Then

$$\int_0^{\pi} |f(t)|^2\, dt \le C \int_0^{\pi} |\phi(t)|^2\, dt = C\, \|\phi(t)\|_{L_{2,\rho}(0,\pi)} .$$

So the operator L_x is invertible in $L_{2,\rho}(0,\pi)$. Then according to Theorem 3 from [26] (see p. 275) it is sufficient to prove that the equation

$$\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) +$$

$$+ \int_0^{\mu^+(x)} A(\xi) F_0(\xi,t)\, d\xi = 0 \tag{3.2}$$

has only trivial solution $A(t) = 0$.

Let $A(t)$ be a non-trivial solution of (3.2). Then

$$\int_0^x \rho(t) \left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) \right)^2 dt +$$

$$+ \int_0^x \rho(t) \left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) \right) \times$$

$$\times \int_0^{\mu^+(x)} A(\xi) F_0(\xi,t)\, d\xi dt = 0.$$

From (2.2) we have

$$\int_0^x \rho(t) \left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) \right)^2 dt +$$

$$+ \int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}} A(\mu^+(t)) \int_0^{\mu^+(x)} A(\xi) \times$$

$$\times \sum_{n=1}^{\infty} \left(\frac{\varphi_0(t,\lambda_n)\cos\lambda_n\xi}{\alpha_n} - \frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0} \right) d\xi dt +$$

$$+ \int_0^x \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) \int_0^{\mu^+(x)} A(\xi) \times$$

$$\times \sum_{n=1}^{\infty} \left(\frac{\varphi_0(t,\lambda_n)\cos\lambda_n\xi}{\alpha_n} - \frac{\varphi_0(t,\lambda_n^0)\cos\lambda_n^0\xi}{\alpha_n^0} \right) d\xi dt = 0.$$

Using (1.7) and (2.9) we obtain

$$\int_0^x \rho(t) \left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) \right)^2 dt +$$

$$+ \int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}} A(\mu^+(t)) \int_0^a A(\xi) \sum_{n=1}^{\infty} \frac{\varphi_0(\xi,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n} d\xi dt +$$

$$+ \int_0^x \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) \int_0^a A(\xi) \sum_{n=1}^{\infty} \frac{\varphi_0(\xi,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n} d\xi dt +$$

$$\begin{gathered}
+\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_a^{\alpha x-\alpha a+a} A(\xi)\times \\
\times\sum_{n=1}^{\infty}\frac{2\alpha}{1+\alpha}\frac{\varphi_0\left(\frac{\xi}{\alpha}+a-\frac{a}{\alpha},\lambda_n\right)\varphi_0\left(t,\lambda_n\right)}{\alpha_n}d\xi dt+ \\
+\int_0^x \frac{1-\sqrt{\rho\left(2a-t\right)}}{1+\sqrt{\rho\left(2a-t\right)}}A(2a-t)\int_a^{\alpha x-\alpha a+a} A(\xi)\times \\
\times\sum_{n=1}^{\infty}\frac{2\alpha}{1+\alpha}\frac{\varphi_0\left(\frac{\xi}{\alpha}+a-\frac{a}{\alpha},\lambda_n\right)\varphi_0\left(t,\lambda_n\right)}{\alpha_n}d\xi dt+ \\
+\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_a^{\alpha x-\alpha a+a} A(\xi)\times \\
\times\sum_{n=1}^{\infty}\frac{1-\alpha}{1+\alpha}\frac{\varphi_0\left(2a-\xi,\lambda_n\right)\varphi_0\left(t,\lambda_n\right)}{\alpha_n}d\xi dt+ \\
+\int_0^x \frac{1-\sqrt{\rho\left(2a-t\right)}}{1+\sqrt{\rho\left(2a-t\right)}}A(2a-t)\int_a^{\alpha x-\alpha a+a} A(\xi)\times \\
\times\sum_{n=1}^{\infty}\frac{1-\alpha}{1+\alpha}\frac{\varphi_0\left(2a-\xi,\lambda_n\right)\varphi_0\left(t,\lambda_n\right)}{\alpha_n}d\xi dt- \\
-\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_0^a A(\xi)\sum_{n=1}^{\infty}\frac{\varphi_0(\xi,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}d\xi dt- \\
-\int_0^x \frac{1-\sqrt{\rho\left(2a-t\right)}}{1+\sqrt{\rho\left(2a-t\right)}}A(2a-t)\int_0^a A(\xi)\sum_{n=1}^{\infty}\frac{\varphi_0(\xi,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}d\xi dt- \\
-\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_a^{\alpha x-\alpha a+a} A(\xi)\times \\
\times\sum_{n=1}^{\infty}\frac{2\alpha}{1+\alpha}\frac{\varphi_0\left(\frac{\xi}{\alpha}+a-\frac{a}{\alpha},\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0}d\xi dt- \\
-\int_0^x \frac{1-\sqrt{\rho\left(2a-t\right)}}{1+\sqrt{\rho\left(2a-t\right)}}A(2a-t)\int_a^{\alpha x-\alpha a+a} A(\xi)\times \\
\times\sum_{n=1}^{\infty}\frac{2\alpha}{1+\alpha}\frac{\varphi_0\left(\frac{\xi}{\alpha}+a-\frac{a}{\alpha},\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0}d\xi dt- \\
-\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_a^{\alpha x-\alpha a+a} A(\xi)\times \\
\times\sum_{n=1}^{\infty}\frac{1-\alpha}{1+\alpha}\frac{\varphi_0\left(2a-\xi,\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0}d\xi dt- \\
-\int_0^x \frac{1-\sqrt{\rho\left(2a-t\right)}}{1+\sqrt{\rho\left(2a-t\right)}}A(2a-t)\int_a^{\alpha x-\alpha a+a} A(\xi)\times \\
\times\sum_{n=1}^{\infty}\frac{1-\alpha}{1+\alpha}\frac{\varphi_0\left(2a-\xi,\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0}d\xi dt=0.
\end{gathered}$$

Substituting $\xi \to \frac{\xi}{\alpha} + a - \frac{a}{\alpha}$ in third, fourth, ninth, and tenth double integrals and $\xi \to 2\alpha - \xi$ in fifth, sixth, eleventh and twelfth double integrals we get

$$\int_0^x \rho(t)\left(\frac{2}{1+\sqrt{\rho(t)}}A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}}A(2a-t)\right)^2 dt+$$

$$+\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_0^a A(\xi)\sum_{n=1}^{\infty}\frac{\varphi_0(\xi,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}d\xi dt+$$

$$+\int_0^x \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}}A(2a-t)\int_0^a A(\xi)\sum_{n=1}^{\infty}\frac{\varphi_0(\xi,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}d\xi dt+$$

$$+\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_a^x A(\mu^+(\xi))\times$$

$$\times\sum_{n=1}^{\infty}\frac{2\alpha^2}{1+\alpha}\frac{\varphi_0(\xi,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}d\xi dt+$$

$$+\int_0^x \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}}A(2a-t)\int_a^x A(\mu^+(\xi))\times$$

$$\times\sum_{n=1}^{\infty}\frac{2\alpha^2}{1+\alpha}\frac{\varphi_0(\xi,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}d\xi dt+$$

$$+\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_{-\alpha x+\alpha a+a}^a A(2a-\xi)\times$$

$$\times\sum_{n=1}^{\infty}\frac{1-\alpha}{1+\alpha}\frac{\varphi_0(\xi,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}d\xi dt+$$

$$+\int_0^x \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}}A(2a-t)\int_{-\alpha x+\alpha a+a}^a A(2a-\xi)\times$$

$$\times\sum_{n=1}^{\infty}\frac{1-\alpha}{1+\alpha}\frac{\varphi_0(\xi,\lambda_n)\varphi_0(t,\lambda_n)}{\alpha_n}d\xi dt-$$

$$-\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_0^a A(\xi)\sum_{n=1}^{\infty}\frac{\varphi_0(\xi,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}d\xi dt-$$

$$-\int_0^x \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}}A(2a-t)\int_0^a A(\xi)\times$$

$$\times\sum_{n=1}^{\infty}\frac{\varphi_0(\xi,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}d\xi dt-$$

$$-\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}}A(\mu^+(t))\int_a^x A(\mu^+(\xi))\times$$

$$\times\sum_{n=1}^{\infty}\frac{2\alpha^2}{1+\alpha}\frac{\varphi_0(\xi,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0}d\xi dt-$$

$$-\int_0^x \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}}A(2a-t)\int_a^x A(\mu^+(\xi))\times$$

$$\times \sum_{n=1}^{\infty} \frac{2\alpha^2}{1+\alpha} \frac{\varphi_0(\xi,\lambda_n^0)\varphi_0(t,\lambda_n^0)}{\alpha_n^0} d\xi dt -$$

$$-\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho\,(t)}} A(\mu^+(t)) \int_{-\alpha x+\alpha a+a}^{a} A(2a-\xi) \times$$

$$\times \sum_{n=1}^{\infty} \frac{1-\alpha}{1+\alpha} \frac{\varphi_0\left(\xi,\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0} d\xi dt -$$

$$-\int_0^x \frac{1-\sqrt{\rho\,(2a-t)}}{1+\sqrt{\rho\,(2a-t)}} A(2a-t) \int_{-\alpha x+\alpha a+a}^{a} A(2a-\xi) \times$$

$$\times \sum_{n=1}^{\infty} \frac{1-\alpha}{1+\alpha} \frac{\varphi_0\left(\xi,\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0} d\xi dt = 0,$$

from which we have

$$\int_0^x \rho(t) \left(\frac{2}{1+\sqrt{\rho\,(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho\,(2a-t)}}{1+\sqrt{\rho\,(2a-t)}} A(2a-t) \right)^2 dt +$$

$$+\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho\,(t)}} A(\mu^+(t)) \int_0^x \frac{2\rho(\xi)}{1+\sqrt{\rho\,(\xi)}} \times$$

$$\times A(\mu^+(\xi)) \sum_{n=1}^{\infty} \frac{\varphi_0\left(\xi,\lambda_n\right)\varphi_0\left(t,\lambda_n\right)}{\alpha_n} d\xi dt +$$

$$+\int_0^x \frac{1-\sqrt{\rho\,(2a-t)}}{1+\sqrt{\rho\,(2a-t)}} A(2a-t) \int_0^x \frac{2\rho(\xi)}{1+\sqrt{\rho\,(\xi)}} \times$$

$$\times A(\mu^+(\xi)) \sum_{n=1}^{\infty} \frac{\varphi_0\left(\xi,\lambda_n\right)\varphi_0\left(t,\lambda_n\right)}{\alpha_n} d\xi dt +$$

$$+\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho\,(t)}} A(\mu^+(t)) \int_0^x \frac{1-\sqrt{\rho\,(2a-\xi)}}{1+\sqrt{\rho\,(2a-\xi)}} \times$$

$$\times A(2a-\xi) \sum_{n=1}^{\infty} \frac{\varphi_0\left(\xi,\lambda_n\right)\varphi_0\left(t,\lambda_n\right)}{\alpha_n} d\xi dt +$$

$$+\int_0^x \frac{1-\sqrt{\rho\,(2a-t)}}{1+\sqrt{\rho\,(2a-t)}} A(2a-t) \int_0^x \frac{1-\sqrt{\rho\,(2a-\xi)}}{1+\sqrt{\rho\,(2a-\xi)}} \times$$

$$\times A(2a-\xi) \sum_{n=1}^{\infty} \frac{\varphi_0\left(\xi,\lambda_n\right)\varphi_0\left(t,\lambda_n\right)}{\alpha_n} d\xi dt -$$

$$-\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho\,(t)}} A(\mu^+(t)) \int_0^x \frac{2\rho(\xi)}{1+\sqrt{\rho\,(\xi)}} \times$$

$$\times A(\mu^+(\xi)) \sum_{n=1}^{\infty} \frac{\varphi_0\left(\xi,\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0} d\xi dt -$$

$$-\int_0^x \frac{1-\sqrt{\rho\,(2a-t)}}{1+\sqrt{\rho\,(2a-t)}} A(2a-t) \int_0^x \frac{2\rho(\xi)}{1+\sqrt{\rho\,(\xi)}} \times$$

$$\times A(\mu^+(\xi)) \sum_{n=1}^{\infty} \frac{\varphi_0\left(\xi,\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0} d\xi dt -$$

$$-\int_0^x \frac{2\rho(t)}{1+\sqrt{\rho(t)}} A(\mu^+(t)) \int_0^x \frac{1-\sqrt{\rho(2a-\xi)}}{1+\sqrt{\rho(2a-\xi)}} \times$$
$$\times A(2a-\xi) \sum_{n=1}^{\infty} \frac{\varphi_0\left(\xi,\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0} d\xi dt -$$
$$-\int_0^x \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) \int_0^x \frac{1-\sqrt{\rho(2a-\xi)}}{1+\sqrt{\rho(2a-\xi)}} \times$$
$$\times A(2a-\xi) \sum_{n=1}^{\infty} \frac{\varphi_0\left(\xi,\lambda_n^0\right)\varphi_0\left(t,\lambda_n^0\right)}{\alpha_n^0} d\xi dt = 0.$$

Thus we obtain

$$\int_0^x \rho(t)\left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t)\right)^2 dt +$$
$$+\sum_{n=1}^{\infty} \frac{1}{\alpha_n}\left(\int_0^x \rho(t)\left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t)\right)\varphi_0(t,\lambda_n)\, dt\right)^2 -$$
$$-\sum_{n=1}^{\infty} \frac{1}{\alpha_n^0}\left(\int_0^x \rho(t)\left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t)\right)\varphi_0\left(t,\lambda_n^0\right) dt\right)^2 = 0.$$

Using the Parseval's equality

$$\int_0^x \rho(t) f^2(t) dt = \sum_{n=1}^{\infty} \frac{1}{\alpha_n^0}\left(\int_0^x \rho(t) f(t) \varphi_0\left(t,\lambda_n^0\right) dt\right)^2$$

for the function

$$f(t) = \frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) \in L_2(0,x)$$

we have

$$\sum_{n=1}^{\infty} \frac{1}{\alpha_n}\left(\int_0^x \rho(t)\left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t)\right)\varphi_0(t,\lambda_n)\, dt\right)^2 = 0$$

or

$$\int_0^x \rho(t)\left(\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t)\right)\varphi_0(t,\lambda_n)\, dt = 0, \quad n \geq 1.$$

Since the system $\{\varphi_0(t,\lambda_n)\}_{n\geq 1}$ is compete in $L_{2,\rho}(0,\pi)$, we have

$$\frac{2}{1+\sqrt{\rho(t)}} A(\mu^+(t)) + \frac{1-\sqrt{\rho(2a-t)}}{1+\sqrt{\rho(2a-t)}} A(2a-t) = 0,$$

i.e. $(L_x A)(t) = 0$, where the operator L_x is defined by (3.1). From invertibility of L_x in $L_{2,\rho}(0,\pi)$ we get $A(x,.) = 0$. □

Theorem 3.2. *Let L and $\widetilde{L}$ be two boundary value problems and*

$$\lambda_n = \widetilde{\lambda}_n, \ \alpha_n = \widetilde{\alpha}_n, \ \ (n \in \mathbf{Z}).$$

Then

$$q(x) = \widetilde{q}(x) \ x \in [0,\pi].$$

Proof. According to (2.2) and (2.3) $F_0(x,t)=\widetilde{F}_0(x,t)$ and $F(x,t)=\widetilde{F}(x,t)$. Then from the main equation (2.1), we have $A(x,t)=\widetilde{A}(x,t)$. It follows from (1.5) that $q(x)=\widetilde{q}(x)$ $x\in[0,\pi]$. □

4. Example

Using [27], we can transform the main equation (2.1) to the following equation:

$$\tilde{A}(x,t)+F(x,t)+\int_0^x \tilde{A}(x,\xi)F(x,\xi)d\xi, \quad 0<t<x, \tag{4.1}$$

where

$$F(x,t)=\rho(t)\sum_{n=1}^{\infty}\left(\frac{\varphi_0(t,\lambda_n)\varphi_0(x,\lambda_n)}{\alpha_n}+\frac{\varphi_0(t,\lambda_n^0)\varphi_0(x,\lambda_n^0)}{\alpha_n^0}\right) \tag{4.2}$$

and

$$\tilde{A}(x,t)=\left\{\begin{array}{c} A(x,t) \quad , \quad 0<t<x \\ A(x,t) \quad , \quad 0<t<-\alpha x+\alpha a+a \\ A(x,t)+\frac{1-\alpha}{1+\alpha}A(x,2a-t) \quad , \quad -\alpha x-\alpha a+a<t<a<x \\ \frac{2\alpha^2}{1+\alpha}A(x,\alpha t-\alpha a+a) \quad , \quad a<t<x. \end{array}\right.$$

We assume that $\lambda_n=\lambda_n^0=\frac{\pi}{\mu^+(\pi)}\left(n-\frac{1}{2}\right), \quad n\geq 1; \quad \alpha_n=\pi, \quad n>1; \quad \alpha_1=\frac{\pi}{2}; \quad \alpha_n^0=\pi, \quad n\geq 1$. From the formula (4.2), we obtain

$$F(x,t)=\frac{1}{\pi}\rho(t)\varphi_0(t,\lambda_1)\varphi_0(x,\lambda_1). \tag{4.3}$$

Substituting (4.3) into the main equation (4.1) we obtain

$$\tilde{A}(x,t)=-\frac{1}{\pi}\rho(t)\varphi_0(t,\lambda_1)\left[\varphi_0(x,\lambda_1)+\int_0^x \tilde{A}(x,\xi)\varphi_0(\xi,\lambda_1)d\xi\right],$$

$$\tilde{A}(x,t)=-\frac{1}{\pi}\rho(t)\varphi_0(t,\lambda_1)L(x), \tag{4.4}$$

where

$$L(x)=\varphi_0(x,\lambda_1)+\int_0^x \tilde{A}(x,\xi)\varphi_0(\xi,\lambda_1)d\xi. \tag{4.5}$$

Substituting (4.4) into (4.5) we obtain

$$L(x)=\varphi_0(x,\lambda_1)-\frac{1}{\pi}L(x)\int_0^x \rho(\xi)\varphi_0^2(\xi,\lambda_1)d\xi.$$

Using the (1.6), we calculate the integral into last equation. Then, we have

$$L(x)=\left\{\begin{array}{c} \varphi_0(x,\lambda_1)-\frac{1}{2\pi}L(x)\left(x+\frac{sin2\lambda_1 x}{2\lambda_1}\right) \quad , \quad 0<x<a \\ \varphi_0(x,\lambda_1)-\frac{1}{2\pi}L(x)\Phi(x,\lambda_1) \quad , \quad a<x<\pi, \end{array}\right. \tag{4.6}$$

where

$$\Phi(x,\lambda_1)=\frac{a}{2}+\frac{sin2\lambda_1 a}{4\lambda_1}+$$
$$+\frac{1}{8}(\alpha+1)^2\left[x-a+\frac{1}{2\lambda_1}\left(\frac{sin2\lambda_1(\alpha x-a\alpha+a)}{\alpha}-sin2\lambda_1 a\right)\right]+$$
$$+\frac{1}{2}(\alpha-1)^2\left[x-a-\frac{1}{2\lambda_1}\left(\frac{sin2\lambda_1(-\alpha x+a\alpha+a)}{\alpha}-sin2\lambda_1(2a-x)\right)\right].$$

From (4.6),

$$(4.7)\qquad L(x)=\begin{cases}\varphi_0(x,\lambda_1)\left[1+\frac{1}{2\pi}\left(x+\frac{sin2\lambda_1 x}{2\lambda_1}\right)\right]^{-1} & ,\quad 0<x<a\\ \varphi_0(x,\lambda_1)\left[1+\frac{1}{\pi}\Phi(x,\lambda_1)\right]^{-1} & ,\quad a<x<\pi,\end{cases}$$

Substituting (4.7) into (4.4) we get
(4.8)

$$\tilde{A}(x,t)=\begin{cases}-\frac{1}{\pi}\varphi_0(t,\lambda_1)\varphi_0(x,\lambda_1)\left[1+\frac{1}{2\pi}\left(x+\frac{sin2\lambda_1 x}{2\lambda_1}\right)\right]^{-1} & ,\quad 0<x<a\\ -\frac{1}{\pi}\rho(t)\varphi_0(t,\lambda_1)\varphi_0(x,\lambda_1)\left[1+\frac{1}{\pi}\Phi(x,lambda_1)\right]^{-1} & ,\quad a<x<\pi,\end{cases}$$

where $0<t<x$ and $\lambda_1=\frac{\pi}{2\mu^+(\pi)}$. Thus, we obtain the solution of main equation (2.1). If we use the formula (1.5) then, we obtain the potential $q(x)$.

Acknowledgements

This work is supported by the Scientific and Technological Research Council of Turkey (TUBITAK).

References

[1] Tikhonov AN, Samarskii AA, Equation of mathematical physics. Dover Books on Physics and Chemistry, Dover, New York, 1990.
[2] Yurko VA, Inverse spectral problems and their applications. Saratov, 2001,497 p. (Russian)
[3] Rasulov ML. Methods of Contour Integration. Series in Applied Mathematics and Mechanics, v.3, North-Holland, Amsterdam, 1967.
[4] Akhtyamov AM. Theory of identification of boundary conditions and its applications. Fizmatlit, Moscow, 2009.
[5] Akhtyamov AM. Mouftakhov AV. Identification of boundary conditions using natural frequencies. Inverse Problems in Science and Engineering, 2004, 12(4), 393–408.
[6] Anderssen RS. The effect of discontinuities in destiny and shear velocity on the asymptotic overtone structure of torsional eigenfrequencies of the earth., Geophys. J. R. Astr. Soc., 1977, 50, 303-309.
[7] Hao DN. Methods for inverse heat conduction problems. Peter Lang Verlag, Frankfurt/Main, Bern, New York, Paris, 1998.
[8] Sedipkov AA. The inverse spectral problem for the Sturm–Liouville operator with discontinuous potential. J. Inverse Ill-Posed Probl., 2012, 20, 139–167.
[9] Marchenko VA. Spektral theory of Sturm- Liouville operators. Naukova Dumka, Kiev, 1977, 331 p. (Russian)
[10] Marchenko VA. Strum-Liouville Operators and Their Aplications. Trans. from the Russian by A. Iacob, Birkhauser Verlag, Basel, Boston, Stuttgard, 1986.
[11] Levitan BM, Sargsjan IS. Sturm- Liouville and Dirac Operators. Kluwer Academic Publishers Group, Dordrecht, 1991.
[12] Levitan BM. Inverse Sturm-Liouville Problems. Translated from the Russian by O. E mov. VNU Science Press BV, Utrecht. 1987.
[13] Freiling G, Yurko V. Inverse Sturm-Liouville problems and their applications. Nova Science Publishers, INC. 2008.
[14] Beals R, Deift P, Tomei C. Direct and inverse scattering on the line. vol.28 of Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, USA, 1988.
[15] Levitan BM, Gasymov MG. Determination of differantial operator by two spektra, Uspekhi mat. Nauk, 1964,v.19 issue 2, pp.3-63. (Russian)
[16] Borg G. Eine Umkehrung der Sturm-Liouvilleschen Eigenwertaufgabe. Acta Math. 78 1946, 1-96.
[17] Poschel J, Trubowitz E. Inverse Spectral Theory. Academic Press, New York, 1987.
[18] Hald OH. Discontinuous inverse eigenvalue problems. Comm. Pure Appl. Math. 1984. 37. 539-577

[19] Carlson R. An inverse spectral problem for Sturm- Liouville operators with discontinuous coefficients. Pro. Amer. Math. Soc. 1994. 120(2). 5-9.

[20] Nabiev AA, Amirov RK. On the boundary value problem for the Sturm-Liouville equation with the discontinuous coefficient. Math. Methods Appl. Sci., 2012, doi:10.1002/mma.2714.

[21] Karahan D, Mamedov KhR. Uniqueness of the solution of the inverse problem for one class of Sturm-Liouville operator, Proceedings of IMM of NAS of Azerbaijan, v. 40, Special Issue, 2014, p. 233-244.

[22] Aliev BA, Yakubov YS. Solvability of boundary value problems for second-order elliptic differential-operator equations with a spectral parameter and with a discontinuous coefficient at the highest derivative. Differential Equations, 2014, 50(4), 464–475.

[23] Altinisik N, Kadakal M, Mukhtarov O. Eigenvalues and eigenfunctions of discontinuous Sturm-Liouville problems with eigenparameter dependent boundary conditions. Acta Math. Hung., 2004, 102(1–2), 159–175.

[24] Akhmedova EN, Huseynov HM. The main equation of the inverse Sturm-Liouville problem with discontinuous coefficients. Proceedings of IMM of NAS of Azerbaijan, v. XXVI (XXXIV), Baku, 2007, pp. 17-32.

[25] Akhmedova EN. On representation of solution of Sturm-Liouville equation with discontinuous coefficients. Proceedings of IMM of NAS of Azerbaijan, v.XVI(XXIV), 2002, p. 5-9.

[26] Lusternik LA, Sobolev VI. Elements of the functional analysis. Moscow, Nauka, 1965, 520 p. (Russian)

[27] Akhmedova EN, Huseynov HM. On inverse problem for Sturm-Liouville operator with discontinuous coefficients. Matematika, YDK 517.984, 2010, p. 3-9. (Russian)

Khanlar Residoglu Mamedov,
Science and Letter Faculty,
Mathematics Department,
Mersin University,
333343, Mersin, Turkey
E-mail address: `hanlar@mersin.edu.tr`

Done Karahan,
Science and Letters Faculty,
Mathematics Department,
Harran University,
Sanliurfa, Turkey
E-mail address: `dkarahan@harran.edu.tr`